arXiv:nlin/0108026v1 [nlin.CD] 15 Aug 2001

The homogeneous balance method, Lax pair, Hirota transformation and a general fifth-order KdV equation

Yang Lei, Zhang Fajiang, Wang Yinghai
Department of physics, Lanzhou University, Gansu 730000, China

In this paper, some notes of the homogeneous balance (HB) method are discussed by a kind of general fifth-order KdV (fKdV) equation. Frist, the auto-Bäcklund transformation and lax represents of the higher-order KdV equation(a specific forms of fKdV equation) are obtained by the HB method. Then, the connection of the Hirota transformation and the HB method is discussed by two specific fKdV equations: the Sawada-Kotera equation and the Lax's equation. At the same time, the solitary wave solution of the general fifth-order KdV equation is obtained.

From 70's, a vast variety of the simple and direct methods to find analytic solutions of PDE have been developed. Recently, the homogeneous balance (HB) method has drawn lots of interests in seeking the solitary wave solution[1] and other kinds of solutions[2], the solitary wave solution of some nonlinear PDEs[3] have been obtained by this method and the relation between the HB method and the CK method or Bäcklund transformation has been discussed[4]. Then how doese the HB method connect with Lax pair and Hirota transformation? In this letter, the relation between the HB method and other methods is discussed by a general fifth-order KdV equation. At the same time, the solitary wave solution of a general KdV equation is obtained.

The general form of the fifth-order KdV is written as

$$u_t + au_{xxxxx} + buu_{xxx} + cu_xu_{xx} + du^2u_x = 0, \tag{1}$$

where a, b, c and d are constants. The equation (1) includes lots of equations which have been studied widely [5], when the special values of a, b, c, d are set. According to the HB method, first, supposing the transformation of u,

$$u = \sum_{i=0}^{N} f^{(i)}(\omega(x,t)), \tag{2}$$

then substituting (2) into (1), and balancing the nonlinear term $buu_{xxx} + cu_xu_{xx} + du^2u_x$ and the linear term au_{xxxxx}, N can be decided to be 2, thus the transformation (2) is order to be

$$u = f^{''}\omega_x^2 + f^{'}\omega_{xx}. \tag{3}$$

Substituting (3) into (1), and collecting all homogeneous terms in partial dericatives of $\omega(x,t)$, we have (because the formula is so long, just one part of it is shown here)

$$(d(f'')^2f^{(3)} + cf^{(3)}f^{(4)} + bf''f^{(5)} + af^{(7)})\omega_x^7 + \ldots\ldots = 0. \tag{4}$$

Setting the coefficient of ω_x^7 in (4) to zero yields an ordinary differential equation for f

$$d(f'')^2f^{(3)} + cf^{(3)}f^{(4)} + bf''f^{(5)} + af^{(7)} = 0. \tag{5}$$

Solving (5), the solution of the equation(5) is

$$f = \alpha \ln \omega, \tag{6}$$

where $\alpha = \frac{6b+3c\pm3\sqrt{(2b+c)^2-40ad}}{d}$, it's obviously that the parameter a, b, c, d need to satisfy $(2b+c)^2 - 40ad > 0$. The solution(6) yields

$$f^{(3)}f^{(4)} = -\frac{\alpha}{60}f^{(7)}, \quad f''f^{(5)} = -\frac{\alpha}{30}f^{(7)}, \quad (f'')^2f^{(3)} = \frac{\alpha}{360}f^{(7)}\cdots\cdots, \tag{7}$$

Substituting (7) into (4), formula (4) can be simplified to a linear polynomial of $f', f'', \ldots$, then setting the coefficients of $f', f'', \ldots$ to zero yields a set of partial differential equations for $\omega(x,t)$,

$$\begin{aligned}
&\omega_{xxt} + a\omega_{7x} = 0,\\
&-2\omega_x\omega_{xt} - \omega_t\omega_{xx} + (c\alpha - 35a)\omega_{xxx}\omega_{4x} + (b\alpha - 21a)\omega_{xx}\omega_{5x} - 7a\omega_x\omega_{6x} = 0,\\
&2\omega_t\omega_x^2 + (210a - 10b\alpha - 3c\alpha + d\alpha^2)\omega_{xx}^2\omega_{xxx} + (140a - 4c\alpha)\omega_x\omega_{xxx}^2 + (210a - 5b\alpha - 3c\alpha)\omega_x\omega_{xx}\omega_{4x}\\
&+(42a - b\alpha)\omega_x^2\omega_{5x} = 0,\\
&(630a - 30b\alpha - 9c\alpha + 3d\alpha^2)\omega_{xx}^3 + (1260a - 30b\alpha - 24c\alpha + 2d\alpha^2)\omega_x\omega_{xx}\omega_{xxx} + (210a - 5b\alpha - 2c\alpha)\omega_x^2\omega_{4x} = 0,\\
&(2520a - 90b\alpha - 42c\alpha + 8d\alpha^2)\omega_{xx}^2 + (840a - 20b\alpha - 14c\alpha + d\alpha^2)\omega_x\omega_{xxx} = 0,
\end{aligned} \tag{8}$$

Where the equation $\omega_{xxt}+a\omega_{7x}=0$ which is from the coefficients of f' is a linear PDE. It's easy to know this equation has the travelling wave solution

$$\omega(x,t)=c_0+c_1e^{\beta(x-vt)}, \tag{9}$$

where c_0, $c_1$and v are arbitrary constants, $\beta=-(\frac{v}{a})^{\frac{1}{4}}$. Substituting formula(9) into the set of equations (8), a set of nonlinear algebraic equations are gotten. Solving the set of nonlinear algebraic equations, the parameter a, b, c, d need to satisfy the formula

$$-4b^3-8b^2c-5bc^2-c^3+40abd+30acd+(10ad-c^2-3bc-2b^2)\sqrt{(2b+c)^2-40ad}=0. \tag{10}$$

Then the exact solitary wave solution of the general fifth KdV equation is obtained,

$$u(x,t)=\frac{3c_0c_1(2b+c\pm\sqrt{(2b+c)^2-40ad})\sqrt{v}e^{(\frac{v}{a})^{\frac{1}{4}}(x-vt)}}{\sqrt{a}d(c_1+c_0e^{(\frac{v}{a})^{\frac{1}{4}}(x-vt)})^2}. \tag{11}$$

In this section, by the HB method, the auto-Bäcklund transformation and lax represents of a special fKdV equation[5] are obtained. The fKdV equation is

$$u_t-\frac{1}{4}u_{xxxxx}-\frac{5}{2}uu_{xxx}-5u_xu_{xx}-\frac{15}{2}u^2u_x=0, \tag{12}$$

where $a=-\frac{1}{4}$, $b=-\frac{5}{2}$, $c=-5$, $d=-\frac{15}{2}$. Base on the transformation (3), ordering

$$u=f^{''}\omega_x^2+f^{'}\omega_{xx}+v, \tag{13}$$

where v is a function of x, t, repeating the step (4)-(7), we yield a set of partial differential equations (14-18) for $\omega(x,t)$ and $v(x,t)$

$$-60vv_x\omega_{xx}+4\omega_{xxt}-10\omega_{xx}v_{xxx}-30v^2\omega_{xxx}-20v_{xx}\omega_{xxx}-20v_x\omega_{4x}-10v\omega_{5x}-\omega_{7x}=0, \tag{14}$$

$$\begin{aligned}&60vv_x\omega_x^2-8\omega_x\omega_{xt}-4\omega_t\omega_{xx}+90v^2\omega_x\omega_{xx}+60\omega_xv_{xx}\omega_{xx}+10\omega_x^2v_{xxx}+80v_x\omega_x\omega_{xxx}\\&-20v\omega_{xx}\omega_{xxx}+50v\omega_x\omega_{4x}-5\omega_{xxx}\omega_{4x}+\omega_{xx}\omega_{5x}+7\omega_x\omega_{6x}=0,\end{aligned} \tag{15}$$

$$\begin{aligned}&4\omega_t\omega_x^2-30v^2\omega_x^3-20\omega_x^3v_{xx}-60v_x\omega_x^2\omega_{xx}+30v\omega_x\omega_{xx}^2-40v\omega_x^2\omega_{xxx}-5\omega_{xx}^2\omega_{xxx}+10\omega_x\omega_{xxx}^2\\&+5\omega_x\omega_{xx}\omega_{4x}-11\omega_x^2\omega_{5x}=0,\end{aligned} \tag{16}$$

$$2v_x\omega_x^3+\omega_{xx}^3-2\omega_x\omega_{xx}\omega_{xxx}+\omega_x^2\omega_{4x}=0, \tag{17}$$

and $v(x,t)$ satisfies the equation

$$v_t-\frac{1}{4}v_{xxxxx}-\frac{5}{2}vv_{xxx}-5v_xv_{xx}-\frac{15}{2}v^2v_x=0, \tag{18}$$

which is the same as the equation (12) of $u(x,t)$. Thus, the set of equation (13-18) constitute an invariant-Bäcklund transformation of the fKdV equation (12). Considering the equation (17), the $v_x(x,t)$ can be solved

$$v_x(x,t)=-\frac{\omega_{xx}^3-2\omega_x\omega_{xx}\omega_{xxx}+\omega_x^2\omega_{xxxx}}{\omega_x^3}, \tag{19}$$

then integrating the formula (19), we have

$$v(x,t)=\frac{1}{2}\frac{\omega_{xx}^2}{\omega_x^2}-\frac{\omega_{xxx}}{\omega_x}+\lambda \tag{20}$$

where λ is a integration constant, here $\frac{1}{2}\frac{\omega_{xx}^2}{\omega_x^2}-\frac{\omega_{xxx}}{\omega_x}$ just is the Schwarzian derivative $S_x\omega(x,t)$, namely $v(x,t)=S_x\omega(x,t)+\lambda$. Substituting the formula(20) and the transformation(13) into the equation(12), the equation(12) can be written to be

$$4\omega_t - \omega_x(S_x\omega(x,t))_{xx} - \frac{3}{2}\omega_x(S_x\omega(x,t))^2 - 10\lambda\omega_x S_x\omega(x,t) - 30\lambda^2 = 0. \tag{21}$$

According the equation(21), which is the form of the Schwarzian derivative of the equation (12), then it's readily to obtain the Lax pair[5]

$$\nu_{xx} = \alpha\nu, \qquad \nu_t = \beta\nu_x + \gamma\nu,$$

where

$$\alpha = \lambda - u, \qquad \beta = \frac{1}{2}u_{xx} + \frac{3}{2}u^2 + 2\lambda u + 4\lambda^2, \qquad \gamma = -\frac{1}{2}\beta_x + \alpha,$$

Applied the HB method, the Schwarzian derivative forms of some nonlinear PDEs can be obtained, and the lax representation of the PDEs can be gotten, such as the KdV, mKdV and the higher-order KdV.

In this section, the connection between Hirota transformation and HB method is discussed. In 1970's, Hirota developed a method to obtain the special solution of nonlinear PDE, namely, Hirota transformation which can transform a nonlinear PDE to a bilinear form. For example, the Hirota transformation $u = 2(\lg f)_{xx}$ can transfer the KdV equation to $[D_x(D_t + D_x^3) + \lambda]f \cdot f = 0$, where D is Hirota's binary operator. It's easy to verify that the Hirota transformation $u = 2(\lg f)_{xx}$ can be obtained from the HB method. Then can the Hirota transformation of the PDEs be obtained from the HB method? In this paper, two special forms of the equation (1) (the Sawada-Kotera equation and the Lax's equation) are discussed. The Sawada-Kotera(SK) equation[5] is

$$u_t + u_{xxxxx} + 15uu_{xxx} + 15u_x u_{xx} + 45u^2 u_x = 0. \tag{22}$$

Substituting $a = 1$, $b = 15$, $c = 15$, $d = 45$ into the formula (6), the transformation (3) can be written to $u = 2(\lg f)_{xx}$, which is the Hirota transformation for the SK equation. By this transformation the SK equation is transformed into bilinear form $[D_x(D_t + D_x^5)]f \cdot f = 0$. The Lax's equation[5] is

$$u_t + u_{xxxxx} + 10uu_{xxx} + 20u_x u_{xx} + 90u^2 u_x = 0. \tag{23}$$

By the Hirota transformation $u = 2(\lg\omega)_{xx}$, the Lax's equation is transformed into bilinear form $[D_x(D_t + D_x^5) - \frac{5}{3}D_t(D_t + D_x^3)]f \cdot f = 0$. Substituting $a = 1$, $b = 10$, $c = 20$, $d = 90$ into the formula (6), the constrain condition $(2b + c)^2 - 40ad > 0$ can't be satisfied. This means the equation (5) has no real solution, and the real transformation (3) is absent. So the HB method is invalid to the Lax's equation, and the Hirota transformation $u = 2(\lg\omega)_{xx}$ can't be obtained by HB method. These results show that the HB method can't be use to obtain the Hirota transformation, even in a kind of nonlinear PDE, but for some famous equations(KdV, mKdV, KP, Bugers and *etc*) the Hirota transformation can be obtained by the HB method.

In summary, a kind of general fifth-order KdV equation, which includes lots of fifth-order KdV [5], is investigated in this paper. The solitary wave solution of the fKdV equation is obtained. The auto-Bäcklund transformation and lax represents(which has been obtain by other methods) of a integrate PDE(the higher-order KdV equation) are obtained by the HB method. And the relation of the Hirota transformation and HB method is discussed.

Acknowledgments: This work was supported by the National Natural Science Foundation of China. The corresponding author email address: yhwang@lzu.edu.cn (Wang Yinghai).

Reference

---

[1] M.Wang, Phys.Lett.A, 199 (1995), 169;
M.Wang, Phys.Lett.A, 213 (1996), 279;
M.Wang, Phys.Lett.A, 216 (1996), 67.
[2] Lei Yang, Zhenggang Zhu, Yinghai Wang, Phys.Lett.A, 260 (1999) 55.
[3] Li Zhibin, M.Wang, J.Phys.A, 26 (1993) 6027;
Engui Fan, Hongqing Zhang, Phys.Lett.A, 245 (1998) 389;
Engui Fan, Hongqing Zhang, Phys.Lett.A, 246 (1998) 403;
Lei Yang, Kongqing Yang, Mathematical Reviews Index: CNO 1698945, (1998);
Zhang Jiefang, Int.J.Theo.Phys, 37 (1998) 2449;
Bishri Abdel-Hamid, Phys.Lett.A, 263 (1999) 338.

[4] Engui Fan, Phys.Lett.A, 265 (2000) 353.
[5] E.A.Jackson, *Perspectives of nonlinear dynamics: Volume 2*, Cambridge University Press, 1990;
K.Sawada, T.Kotera, Prog.Theor.Phys, 51 (1974) 1355;
A.P.Fordy, J.Gibbons, J.Math.Phys, 22 (1981) 1170;
P.J.Caudrey, R.K.Dodd, J.D.Gibbon, Proc.r.Soc.A, 351 (1976) 407;
H.Morris, J.Math.Phys, 18 (1977) 530;
P.G.Drazin, R.S.Johnson, *Solitons: an Introduction*, Cambridge University Press, 1989;
A.P.Fordy, J.Gibbons, J.Math.Phys, 21 (1980) 2508;
M.Ito, J.Phys.Soc.Jpn, 49 (1980) 771.